

Dr. Martin J.R.R. Vanlierde

B.A., B.Sc., MB.ChB. (U.C.T.), L.M.C.C., FRCS (Edin.), FCS (SA) Otol.

OTORHINOLARYNGOLOGIST

Pr No: 3003612

Sinonasal Surgery

Cosmetic Rhinoplasty & Botox Therapy

Laryngology

Snoring Surgery

General ENT

Endoscopic DCR

Head & Neck Surgery

1002 Vincent Pallotti Hospital

Alexandra Road

Pinelands, 7405

Tel: 021 – 5322156

Fax: 021 – 5323876

Email: vlrecep1@surgeon.co.za

175 Milnerton Medi-Clinic

Racecourse Road,

Milnerton, 7441

Tel: 021 – 5517211

Fax: 021 – 5517230

Email: vlrecep2@surgeon.co.za

POST-TONSILLECTOMY INSTRUCTIONS

1. The patient must stay in bed on the day of discharge from hospital.
2. The patient may be allowed up (indoors) the day following the operation.
3. The patient may be allowed outside the following day, but must remain indoors at night for 10 days and off from work / school for 7 days.
4. The pain after tonsillectomy is due to stiffness of the throat muscles. The more the patient chews and eats, the sooner the pain will be relieved. Encourage eating of solid foods, e.g. Bread and a full diet should be started as soon as possible. Biltong is a good food to chew. Panado or another analgesic thirty minutes before a meal will relieve the pain and allow solid foods to be eaten. Avoid eating bananas, acid or salty foods and condiments.
5. During the first post-operative week a sore throat, earache and a slight rise in temperature are normal responses and should not cause concern. If however these symptoms are severe or if bleeding occurs, or if you are concerned in any way, please contact either your local GP or myself.
6. The area from which the tonsils were removed will heal with the formation of a yellow scab. This is normal healing pattern, and does not indicate infection.

Please expect a period of tiredness and irritability after surgery. Whatever the effort made, having an operation is still a very stressful experience and rest, good food, lots of fluids to drink and plenty of sleep will speed you towards recovery!